



## Cat nr AE00154

## **Product Datasheet**

Mouse Monoclonal Antibody, clone ADFP/1365 to:

## PLIN2, Perilipin-2

Product performance

Adipophilin; Adipose differentiation-related protein; Perilipin-2; ADFP; ADRP


ER, plasma membrane, cytoplasmic lipid droplets.
PLIN2 123 Q99541
PA, WB HepG2, JAR, liver, adrenal gland, cerebellum. 85
By Protein G from bioreactor concentrate 200ug IgG/ml in PBS, 0.05% BSA, 0.05% azide (20ug or 100ug) 1mg IgG/ml in PBS (100ug or contact us for quotation) 20ug 100ug Mouse IgG1, kappa Human Recombinant fragment around aa 249-376 of human PLIN2 protein (exact sequence is proprietary) Within aa 249-376 region
Avoid repeated freeze/thaw cycles. For long term storage, keep small aliquots at -20C or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve
antibody at 4C for 6-12 months, when kept away from direct sun light.  Integrity warranted for 24 months after purchase when handled and stored
according to instructions, see below.  This product is only warranted for the specifications as described in this product sheet and only when the product is handled and stored according to instructions. User should validate this antibody in the application and tissue/cell type as required, after confirmation of integrity upon receipt is obtained by reproducing the performance as described below. Should such confirmation not be attempted, any warranty is void. In case of non-conformance, user needs to contact us immediately for replacement or refund.
This product is for in vitro research use only. Any other applications, such as diagnostics or therapeutics, or in vivo experiments, and the validation of this product therein, are solely at the responsibility of the buyer/user.

see next pages

## **Product data:**


Western Blot (WB):

This product was successfully used to stain an approx. 48kDa band in lysates of cell line Jurkat. Recommended concentration: 1-3ug/ml




Western Blot of a Jurkat lysate (30ug) stained with PLIN2 Mouse Monoclonal Antibody AE00154 at 1ug/ml (1h at ambient temp). ECL staining by HRP.

SDS-PAGE Analysis of Purified PLIN2 Mouse Monoclonal Antibody AE00154. Confirmation of Purity and Integrity of Antibody.



Integrity of the purified antibody AE00154 under non-reduced and reduced conditions, showing intact IgG at around 160kDa (NR) and intact heavy and light chains at 50kDa and 25kDa resp. (R).

Specificity and selectivity of AE00154 were tested against >19,000 full-length human proteins on a human protein array. A protein BLAST search against H. sapiens revealed the following closely related protein: PLIN1, 3 and 4. The proteins PLIN1 and 3 were part of the array used and showed no cross-reactivity signals.



Cross-reactivity assessment of PLIN2 Mouse Monoclonal Antibody AE00154 (1ug/ml) on CDI's Protein Array containing more than 19,000 full-length human proteins.

The Z-score represents the strength of a signal that an antibody (through a fluorophore-tagged secondary reagent) produces when binding to a particular protein on the array. Z-scores are in units of standard deviations (SD's) above the mean value of all signals generated on that array. When Z-scores are arranged in descending order, the difference between two successive values will be the S-score for the first. Thus, the S-score represents the relative specificity of the antibody to its intended target. An antibody is considered specific to its intended target, when it has an S-score of at least 2.5. For example, if an antibody binds to intended protein X with a Z-score of 43 and to the cross-reacting protein Y with a next Z-score of 14, then the S-score for the antibody to intended target X equals 29 (43-14).