



Cat nr AE00212

Product Datasheet

Rabbit Polyclonal Phospho-Specific Antibody, EGFR-pY845 to: **EGFR**, phosphorylated Y845


Epithelial Growth Factor Receptor; EGFR; ERBB; ERBB1; HER1; NISBD2; PIG61; mENA

Cellular localization Official Symbol (Gene) GeneID SwissProt	Plasma membrane, cell surface. Translocation to nucleus upon EGF stimulation. EGFR 1956 P00533
Confirmed Applications Positive controls Customers positive feedback Aeonian Rating®	WB EGF-stimulated serum-starved A431 70
Purification	Antigen-affinity purified from serum by sequential phospho- peptide and non-phospho-peptide affinity columns PBS, 0.02% azide, 50% glycerol
Amount Isotype Confirmed species reactivity Immunogen	 100ug 200ug Rabbit IgG Human A synthetic peptide around phosphorylated Y869 of human EGFR (NP_005219.2) corresponding to Y845 of the mature protein (starts with L25).
Epitope	A short sequence around phosphorylated Y845 (exact sequence is proprietary)
Storage instructions	Avoid repeated freeze/thaw cycles. For long term storage, keep small aliquots at -20C or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve antibody at 4C for 6-12 months, when kept away from direct sun light.
Expiration	Integrity warranted for 24 months after purchase when handled and stored according to instructions, see below.
Warranty	This product is only warranted for the specifications as described in this product sheet and only when the product is handled and stored according to instructions. User should validate this antibody in the application and tissue/cell type as required, after confirmation of integrity upon receipt is obtained by reproducing the performance as described below. Should such confirmation not be attempted, any warranty is void. In case of non-conformance, user needs to contact us immediately for replacement or refund.
Liability	This product is for in vitro research use only. Any other applications, such as diagnostics or therapeutics, or in vivo experiments, and the validation of this product therein, are solely at the responsibility of the buyer/user.

Product data:

Western Blot (WB):

This product was successfully used to stain an approx. 200kDa band in lysates of A431 after overnight serum starvation and subsequent treatment with EGF. Recommended concentration: 1-3ug/ml



Western Blot of 30ug lysate from serum-starved A431 with or without stimulation by 100mg/ml EGF (30 min, 37C)), stained with EGFR-pY845 Rabbit Polyclonal Antibody AE00212 at 1ug/ml (1h at ambient temp). ECL staining by HRP.