



## Cat nr AE00275

Product performance

## **Product Datasheet**

Mouse Monoclonal Antibody, clone GH/3155 to:

## **GH1,** Growth Hormone 1

Growth Hormone; Pituitary growth hormone; Somatotropin; GH; GH1; GH-N; GHB5; GHN; IGHD1B; hGH-N


Cellular localization	Cytoplasm and secreted
Official Symbol (Gene) GeneID SwissProt	GH1 2688 P01241
Confirmed Applications Positive controls	IHC, PA Pituitary gland
Aeonian Rating©	80
Purification Formulation  Amount Isotype Confirmed species reactivity Immunogen	By Protein G from bioreactor concentrate  200 ug IgG/ml in PBS, 0.05% BSA, 0.05% azide (20 ug or 100 ug)  1mg IgG/ml in PBS (100 ug or contact us for quotation)  20 ug 100 ug  Mouse IgG1, kappa  Human  Full length of the human GH1 protein
Epitope	Unknown
Storage instructions	Avoid repeated freeze/thaw cycles. For long term storage, keep small aliquots at -20C or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve antibody at 4C for 6-12 months, when kept away from direct sun light.
Storage instructions  Expiration	or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve
	or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve antibody at 4C for 6-12 months, when kept away from direct sun light.  Integrity warranted for 24 months after purchase when handled and stored
Expiration	or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve antibody at 4C for 6-12 months, when kept away from direct sun light.  Integrity warranted for 24 months after purchase when handled and stored according to instructions, see below.  This product is only warranted for the specifications as described in this product sheet and only when the product is handled and stored according to instructions.  User should validate this antibody in the application and tissue/cell type as required, after confirmation of integrity upon receipt is obtained by reproducing the performance as described below. Should such confirmation not be attempted, any warranty is void. In case of non-conformance, user needs to contact us immediately

see next pages

## **Product data:**


ImmunoHistoChemistry (IHC):

This product shows staining of somatotrophs in human anterior pituitary sections. Recommended concentration: 1-3ug/ml




Formaldehyde-fixed, paraffin-embedded human anterior pituitary stained with GH1 Mouse Monoclonal Antibody AE00275 at 1-2ug/ml for 30 minutes at RT. Epitope retrieval: Boiling at pH6 for 10-20 min followed by 20 min cooling. DAB staining by HRP polymer.

SDS-PAGE Analysis of Purified GH1 Mouse Monoclonal Antibody AE00275. Confirmation of Purity and Integrity of Antibody.



Integrity of the purified antibody AE00275 under non-reduced and reduced conditions, showing intact IgG at around 140kDa (NR) and intact heavy and light chains at 45kDa and 25kDa resp. (R).

Specificity and selectivity of AE00275 to GH1 were tested against >19,000 full-length human proteins on a human protein array. A protein BLAST search against H. sapiens revealed the following closely related proteins: CSH1, CSHL1 and CSH2. These proteins were part of the array used and showed no cross-reactivity signals.



Cross-reactivity assessment of GH1 Mouse Monoclonal Antibody AE00275 (lug/ml) on CDI's Protein Array containing more than 19,000 full-length human proteins.

The Z-score represents the strength of a signal that an antibody (through a fluorophore-tagged secondary reagent) produces when binding to a particular protein on the array. Z-scores are in units of standard deviations (SD's) above the mean value of all signals generated on that array. When Z-scores are arranged in descending order, the difference between two successive values will be the S-score for the first. Thus, the S-score represents the relative specificity of the antibody to its intended target. An antibody is considered specific to its intended target, when it has an S-score of at least 2.5. For example, if an antibody binds to intended protein X with a Z-score of 43 and to the cross-reacting protein Y with a next Z-score of 14, then the S-score for the antibody to intended target X equals 29 (43-14).