



## Cat nr AE00242

## **Product Datasheet**

Recombinant Rabbit Antibody, clone C1A/1506R to:

## CD1a

CD1a molecule; CD\_antigen: CD1a; T-cell surface glycoprotein CD1a; T-cell surface antigen T6/Leu-6; hTa1 thymocyte antigen; CD1; FCB6; HTA1; R4; T6


Cellular localization	Plasma membrane, cell surface, endosomes		
Official Symbol (Gene)	CD1A		
GeneID	909		
SwissProt	P06126		
Confirmed Applications	IHC, PA		
Positive controls	skin		
Aeonian Rating©	90		
Purification	By Protein A from bioreactor concentrate		
Formulation	200ug IgG/ml in PBS, 0.05% BSA, 0.05% azide (20ug or 100ug)		
Amount	1mg lgG/ml in PBS (100ug or contact us for quotation)  20ug 100ug		
Isotype	Rabbit IgG		
Confirmed species reactivity	Human		
Immunogen	Recombinant full-length human CD1a protein		
mmunogen	Recombinant fair feilgen naman es 1a procein		
Epitope	Unknown		
Storage instructions	Avoid repeated freeze/thaw cycles. For long term storage, keep small aliquots at -20C		
Storage mistractions	or -80C and keep one aliquot at 4C for daily experimentations. Azide will preserve antibody at 4C for 6-12 months, when kept away from direct sun light.		
Expiration	Integrity warranted for 24 months after purchase when handled and stored according to instructions, see below.		
Warranty	This product is only warranted for the specifications as described in this product sheet and only when the product is handled and stored according to instructions. User should validate this antibody in the application and tissue/cell type as required, after confirmation of integrity upon receipt is obtained by reproducing the performance as described below. Should such confirmation not be attempted, any warranty is void. In case of non-conformance, user needs to contact us immediately for replacement or refund.		
Liability	This product is for in vitro research use only. Any other applications, such as diagnostics or therapeutics, or in vivo experiments, and the validation of this product therein, are solely at the responsibility of the buyer/user.		
Product performance	see next pages		

## **Product data:**

ImmunoHistoChemistry (IHC):


This product was successfully used to stain Langerhans cells in human skin sections. Recommended  $\,$ 

concentration: 1-3ug/ml




Formaldehyde-fixed, paraffin-embedded human skin stained with CD1a Rabbit Recombinant Antibody AE00242 at 1-2ug/ml for 30 minutes at RT. Epitope retrieval: boiling at pH6 for 10-20 min followed by 20 min cooling. DAB staining by HRP polymer.

SDS-PAGE Analysis of Purified CD1a Recombinant Rabbit Antibody AE00242. Confirmation of Purity and Integrity of Antibody.



Integrity of the purified antibody AE00242 under non-reduced and reduced conditions, showing intact IgG at around 120kDa (NR) and intact heavy and light chains at 50kDa and 25kDa resp. (R).

Specificity and selectivity of AE00242 to CD1a were tested against >19,000 full-length human proteins on a human protein array. A protein BLAST search against H. sapiens revealed the following closely related other proteins: CD1b, c, d and e. The related proteins CD1b, c and e were part of the array used and showed no cross-reactivity signals.



Cross-reactivity assessment of CD1a Rabbit Recombinant Antibody AE00242 (1ug/ml) on CDI's Protein Array containing more than 19,000 full-length human proteins.

The Z-score represents the strength of a signal that an antibody (through a fluorophore-tagged secondary reagent) produces when binding to a particular protein on the array. Z-scores are in units of standard deviations (SD's) above the mean value of all signals generated on that array. When Z-scores are arranged in descending order, the difference between two successive values will be the S-score for the first. Thus, the S-score represents the relative specificity of the antibody to its intended target. An antibody is considered specific to its intended target, when it has an S-score of at least 2.5. For example, if an antibody binds to intended protein X with a Z-score of 43 and to the cross-reacting protein Y with a next Z-score of 14, then the S-score for the antibody to intended target X equals 29 (43-14).